

# Creeping Sedge



Carex chordorrhiza



Photo credits: Troy Weldy

**Scientific Name** *Carex chordorrhiza*  
Ehrh. ex L. f.

**Family Name** Cyperaceae  
Sedge Family

## Did you know?

The name chordorrhiza means with cord-like roots (Fernald 1970). This is probably in reference to the vegetative stems which become prostrate and root at the nodes. As they become buried by mosses and other plants they appear like rhizomes (Reznicek and Catling 2002).

## Summary

**Protection** Threatened in New York State, not listed federally.

This level of state protection means: listed species are those with: 1) 6 to fewer than 20 extant sites, or 2) 1,000 to fewer than 3,000 individuals, or 3) restricted to not less than 4 or more than 7 U.S.G.S. 7 ½ minute topographical maps, or 4) listed as threatened by U.S. Department of Interior.

**Rarity** G5, S2

A global rarity rank of G5 means: This species is demonstrably secure globally, though it may be quite rare in parts of its range, especially at the periphery.

A state rarity rank of S2 means: This plant is threatened/imperiled in New York because of rarity (typically 6-20 populations or few remaining individuals) or is vulnerable to extirpation from New York due to biological factors.

## Conservation Status in New York

There are at least fourteen known populations and equally as many historical locations. Most of these are located within Oswego County where many peatland habitats exist. Other populations are present in Warren, Wyoming, and Cattaraugus Counties. This plant is typically found in higher quality inland poor fens and similar habitats. These are relatively well protected, although succession is a possible threat. Many of the populations are large, including some with upwards of 10,000 culms. This sedge is relatively easy to identify, therefore it is not overlooked often.

## Short-term Trends

There are about a dozen populations that have been seen in recent years. Exact trends at these populations are not clear although there is no indication that these populations are declining. At least half of these populations are large and robust. The other known extant populations are smaller. Overall, short term trends are not clear but may indicate that this species is currently stable.

## Long-term Trends

There are at least two populations that appear to have been extirpated sometime in the past century. There are about another dozen populations that have not been seen in recent years. Many of these have very poor locality information. Therefore, finding these historical populations is difficult and it is unknown if these populations are still extant. Overall, long term trends indicate at least some decline.

# Conservation and Management

## Threats

*Phragmites australis* is becoming a serious threat to the habitat at one population. The hydrology at two populations appears to have been altered and perhaps is threatening the habitat where *C. chordorrhiza* occurs. Potential threats include development especially on the edges of the wetlands, trampling by scientists and people fishing, and alteration of the hydrology.

## Conservation Strategies and Management Practices

*Phragmites* should be controlled at one population where it is encroaching the habitat where *Carex chordorrhiza* occurs. Further hydrological changes at or near habitat where *C. chordorrhiza* occurs should be prevented. Where possible, the hydrology of the general area should be restored to pre-altered conditions. In some cases, this is as simple as maintaining culverts.

## Research Needs

Survey work is needed on all historical population.

## Habitat

*Carex chordorrhiza* occurs primarily in peat lands that often are at least somewhat minerotrophic including rich, medium, and poor fens. The habitat is usually open although it does occur where there are some shrubs present and also in openings in forested wetlands (New York Natural Heritage Program 2006). Fens, bogs, floating mats on lakeshores, emergent sedge marshes, usually in very wet sites, often in shallow water (Reznicek and Catling 2002). Sphagnum bogs (Gleason and Cronquist 1991). Quagmires and inundated bogs (Fernald 1970). Found in very wet sphagnum bogs and lake-borders in calcareous districts (Mackenzie 1931-1935).

## Associated Ecological Communities

### Dwarf Shrub Bog

A wetland usually fed by rainwater or mineral-poor groundwater and dominated by short, evergreen shrubs and peat mosses. The surface of the peatland is usually hummocky, with shrubs more common on the hummocks and peat moss throughout. The water in the bog is usually nutrient-poor and acidic.

### Inland Poor Fen

A wetland fed by acidic water from springs and seeps. Plant remains in these fens do not decompose rapidly and thus the plants in these fens usually grow on older, undecomposed plant parts of mostly sphagnum mosses.

### Medium Fen

A wetland fed by water from springs and seeps. These waters are slightly acidic (pH values generally range from 4.5 to 6.5) and contain some dissolved minerals. Plant remains in these fens do not decompose rapidly and thus the plants in these fens usually grow on older, undecomposed plant parts of woody material, grasses, and mosses.

### Rich Shrub Fen

A wetland with many shrubs that is usually fed by water from springs and seeps. These waters have high concentrations of minerals and high pH values, generally from 6.0 to 7.8. Plant remains in these fens do not decompose rapidly and thus the plants in these fens usually grow on older, undecomposed woody plant parts.

## Associated Species

Bog Rosemary (*Andromeda polifolia*)  
American Woollyfruit Sedge (*Carex lasiocarpa*)  
Mud Sedge (*Carex limosa*)  
Leatherleaf (*Chamaedaphne calyculata*)  
Twig Rush (*Cladium mariscoides*)  
Marsh Willow-herb (*Epilobium palustre*)  
Bog Buckbean (*Menyanthes trifoliata*)  
Sweet Bayberry (*Myrica gale*)  
Bog Willow (*Salix pedicellaris*)  
Large Cranberry (*Vaccinium macrocarpon*)

## Identification Comments


Creeping sedge is a grass-like perennial that grows in patches. The leaves are strap-like and 0.4-3.0 mm long. The stems arise singly and are 5-35 cm long. Some of the stems do not produce flowers and these stems fall down and elongate up to 120 cm long. Some of the stems have 2-7 flower/fruit clusters (spikes) towards their apices. These spikes are egg shaped, attach directly to the stems, and are composed of male flowers above and female flowers below. The female flowers mature into fruits (perigynia) which are 2.0-4.5 mm long and have a small beak at their apex (Reznicek and Catling 2002).

## Best Life Stage for Identifying This Species

This species is easiest to identify when it has mature or just immature perigynia. It can be identified at other times of the year based on its vegetative characteristics but not as easily or definitively.

## The Best Time to See

*Carex chordorrhiza* starts to produce immature perigynia in June. These mature and persist till late July or early August. Towards the end of this season the perigynia start to shed easily. This species can be recognized at other times of the year based on vegetative characteristics but it is most obvious when it is in fruit. Therefore, the best time to survey for this species is from the second week in June through mid-July.



**The time of year you would expect to find Creeping Sedge in New York.**

## Similar Species

*Carex chordorrhiza* is a very distinctive sedge especially because it has vegetative culms that root at the nodes. This character is shared by only a few other species of *Carex* which are all quite distinct from *C. chordorrhiza*.

## Conservation Comments

*Carex chordorrhiza* is the only member of section *Chordorrhizae* (Reznicek and Catling 2002).

## Taxonomy

Kingdom Plantae

└ Phylum Anthophyta

└ Class Monocots (Monocotyledoneae)

└ Order Cyperales

└ Family Cyperaceae (Sedge Family)

## Additional Common Names

Sedge

# Additional Resources

## Links

### Identification on YouTube

<http://www.youtube.com/watch?v=afCG6TkTDBc&NR=1>

### New York Flora Atlas

<http://www.newyork.plantatlas.usf.edu/Plant.aspx?id=962>

### USDA Plants Database

<http://plants.usda.gov/java/nameSearch?mode=sciname&keywordquery=CAREX+CHORDORRHIZA>

### NatureServe Explorer

<http://natureserve.org/explorer/servlet/NatureServe?searchName=CAREX+CHORDORRHIZA>

### Google Images

<http://images.google.com/images?q=CAREX+CHORDORRHIZA>

### Flora of North America

[http://efloras.org/florataxon.aspx?flora\\_id=1&taxon\\_id=242357115](http://efloras.org/florataxon.aspx?flora_id=1&taxon_id=242357115)

### Ecology on YouTube

<http://www.youtube.com/watch?v=kymVBr8X-BM>

## Best Identification Reference

Gleason, Henry A. and A. Cronquist. 1991. Manual of Vascular Plants of Northeastern United States and Adjacent Canada. The New York Botanical Garden, Bronx, New York. 910 pp.

## References

Fernald, M.L. 1950. Gray's manual of botany. 8th edition. D. Van Nostrand, New York. 1632 pp.

Holmgren, Noel. 1998. The Illustrated Companion to Gleason and Cronquist's Manual. Illustrations of the Vascular Plants of Northeastern United States and Adjacent Canada. The New York Botanical Garden, Bronx, New York.

Keys, Jr., J.; Carpenter, C.; Hooks, S.; Koenig, F.; McNab, W.H.; Russell, W.; Smith, M.L. 1995. Ecological units of the eastern United States - first approximation (cd-rom), Atlanta, GA: U.S. Department of Agriculture, Forest Service. GIS coverage in ARC/INFO format, selected imagery, and map unit tables.

Mackenzie, K.K. 1931-1935. Cariceae. North American Flora 18: 1-478.

NatureServe. 2005. NatureServe Central Databases. Arlington, Virginia. USA

New York Natural Heritage Program. 2010. Biotics database. New York Natural Heritage Program. New York State Department of Environmental Conservation. Albany, NY.

Reschke, Carol. 1990. Ecological communities of New York State. New York Natural Heritage Program, New York State Department of Environmental Conservation. Latham, NY. 96 pp. plus xi.

Reznicek, A.A. and P.M. Catling. 2002. Carex Linnaeus sect. Chordorrhizae (Heuffel) Meinshausen. Pages 298-299 in Flora of North America Editorial Committee (editors), Flora of North America, North of Mexico, Volume 23, Magnoliophyta: Commelinidae (in part): Cyperaceae. Oxford University Press, New York, NY, USA. 608pp + xxiv.

Weldy, T. and D. Werier. 2010. New York flora atlas. [S.M. Landry, K.N. Campbell, and L.D. Mabe

(original application development), Florida Center for Community Design and Research <http://www.fccdr.usf.edu/>. University of South Florida <http://www.usf.edu/>]. New York Flora Association <http://www.nyflora.org/>, Albany, New York

---

**New York Natural Heritage Program**

625 Broadway, 5th Floor,  
Albany, NY 12233-4757  
Phone: (518) 402-8935  
[acris@nynhp.org](mailto:acris@nynhp.org)

This project is made possible with funding from:

- New York State Department of Environmental Conservation Hudson River Estuary Program
- Division of Lands & Forests, Department of Environmental Conservation
- New York State Office of Parks, Recreation and Historic Preservation

Information for this guide was last updated on Aug 22, 2017

This guide was authored by